

Neural Network identification of halo white dwarfs

Santiago Torres¹, Enrique García-Berro², and Jordi Isern³

Received _____; accepted _____

arXiv:astro-ph/9809363v1 28 Sep 1998

¹Departament de Telecomunicació i Arquitectura de Computadors, EUP de Mataró, Universitat Politècnica de Catalunya, Avda. Puig Cadafalch 101, 08303 Mataró, Spain

²Departament de Física Aplicada, Universitat Politècnica de Catalunya & Institute for Space Studies of Catalonia–UPC, Jordi Girona Salgado s/n, Mòdul B-5, Campus Nord, 08034 Barcelona, Spain

³Institute for Space Studies of Catalonia–CSIC, Edifici Nexus–104, Gran Capità 2–4, 08034 Barcelona, Spain

ABSTRACT

The white dwarf luminosity function has proven to be an excellent tool to study some properties of the galactic disk such as its age and the past history of the local star formation rate. The existence of an observational luminosity function for halo white dwarfs could provide valuable information about its age, the time that the star formation rate lasted, and could also constrain the shape of the allowed Initial Mass Functions (IMF). However, the main problem is the scarce number of white dwarfs already identified as halo stars. In this *Letter* we show how an artificial intelligence algorithm can be successfully used to classify the population of spectroscopically identified white dwarfs allowing us to identify several potential halo white dwarfs and to improve the significance of its luminosity function.

Subject headings: stars: white dwarfs – stars: luminosity function – Galaxy: stellar content

1. Introduction

Halo white dwarfs have received a continuous interest during almost one decade from the theoretical (Mochkovitch et al. 1990; Tamanaha et al. 1990) and observational points of view. From this last point of view Liebert, Dahn & Monet (1989) studied a high proper motion sample and from it derived the first — and up to now the only one available — halo white dwarf luminosity function. Later Flynn, Gould & Bahcall (1996) and Méndez et al. (1996) studied the white dwarf content of the Hubble Deep Field, after the suggestion of the MACHO team that most of the dark matter in the galactic halo could be in the form of white dwarfs (Alcock et al. 1997), with negative results.

The observational white dwarf luminosity function of the halo was obtained from a sample of white dwarfs with known parallaxes, large proper motions ($2.5'' \text{ yr}^{-1} \geq \mu \geq 0.8'' \text{ yr}^{-1}$), using a limiting magnitude of $m_V = 19^{\text{mag}}$ and assuming that only white dwarfs with tangential velocities in excess of 250 km s^{-1} were halo members (Liebert et al. 1989). Consequently only 5 white dwarfs contribute to the luminosity function and, thus, the statistics is very poor. Besides, in the sample of Liebert et al. (1989) there is not any bright halo white dwarf. The absence of bright halo white dwarfs in this sample could be due to an observational bias but this issue, which bears important consequences regarding the IMF of halo stars, remains to be studied, since the biased IMFs recently proposed by Adams & Laughlin (1996) and by Chabrier, Ségreain & Méra (1996) predict very few bright halo white dwarfs if the halo age is large enough (Isern et al. 1998).

In this *Letter* we address the issue of whether or not there exist other halo white dwarfs in the existing catalogs and how do we identify them. For this purpose we use a neural network technique. In the end, this will allow us to present a preliminary luminosity function of halo white dwarfs and compare it with the theoretical predictions.

2. Method and results

With the advent of large astronomical databases the need of efficient techniques to improve automatic classification strategies has lead to a considerable amount of new developments in the field. Among these techniques the most promising ones are based in artificial intelligence algorithms. Neural networks have been used successfully in several fields such as pattern recognition, financial analysis, biology — see Kohonen (1990) for an excellent review — and in astronomy. For instance, Bezell & Peng (1998) used these techniques to automatically discriminate stars from galaxies, Naim et al. (1995) used them to classify galaxies according to their morphology, Serra-Ricart et al. (1996) found the fraction of binaries in stars clusters, and Hernández-Pajares & Floris (1994) used such techniques to classify populations in the *Hipparcos Input Catalogue*.

The common characteristic of all the existing neural network classification techniques is the existence of a learning process very much in the same manner as human experts manually classify. Generally speaking there are two different approaches: the supervised and the unsupervised learning methods. The main advantage of the last class of methods is that require minimum manipulation of the input data and, thus, the results are supposedly more reliable. Their leading exponent is the so-called Kohonen Self-Organizing Map (SOM). A thorough description of this technique is out of the scope of this paper. Therefore, we refer the reader to the specific literature (Kohonen 1997). However for our purpose it is convenient to summarize its basic principle and properties. The basic principle is to map a multi-dimensional input space (S) into a bi-dimensional space (Λ). Similar objects in S (groups) are mapped in nodes in Λ . The most noticeable property of this procedure is the reduction of the dimensionality of the input space allowing, at the same time, the identification of groups in the input data and the automatic classification of individual objects. Besides, neighbor groups in Λ have similar properties.

The catalog of McCook & Sion (1987) compiles the observational data of 1279 white dwarfs. In order to classify the stellar populations presumably present in this catalog a set of variables describing their properties should be adopted. It should be noted that the larger the set of variables adopted, the smaller the number of objects that will have determinations for all the variables. Conversely, if the number of variables in the set is small we could be disregarding valuable information. We have adopted a minimal set in order to be able to analyse the largest possible number of objects in the catalog. The variables adopted in this study are: the absolute visual magnitude M_V , the proper motion μ , the galactic coordinates (l, b) , the parallax π , and a color index, $B - V$. This reduces considerably the number of objects with *all* the determinations, but allows a secure classification. We have found very convenient to use the reduced proper motion defined as $H = M_V - 5 \log \pi + 5 \log \mu$, instead of μ itself because the resulting groups are easier to visualize.

The statistical classification of an observational database usually ends up with the detection of groups in the input space that require an “a posteriori” analysis. Since we are interested in detecting different stellar populations, simultaneously with the clustering process we mix in the input data a synthetic population of tracer stars that will allow us to *label* the groups detected by the classification algorithm as halo, disk or intermediate population. The results of the classification procedure are not sensitive to the fine details of these synthetic populations, except to the IMF. These synthetic tracer stars have been produced using a Monte Carlo (MC) simulator. The description of the MC simulation of the disk population can be found in García-Berro & Torres (1997) or García-Berro et al. (1998). A comprehensive discussion of the results of the MC simulation of the halo population will be published elsewhere. However, and for the sake of completeness, a brief summary of the inputs is given here. We have adopted a standard, Salpeter-like, IMF (Salpeter 1961). The halo was supposed to be formed 14 Gyr ago in an intense burst of star formation of 1 Gyr of duration. The stars are randomly distributed in a sphere of radius 200 pc centered in

the sun according to a density profile given by the expression $\rho(r) \propto (a^2 + R_{\odot}^2)/(a^2 + r^2)$, being r the galactocentric radius, $a \approx 5$ kpc and $R_{\odot} = 8.5$ kpc. The velocities of the tracer stars were randomly drawn according to normal distributions for both the radial and the tangential components, with velocity dispersions as given in Marković & Sommer-Larsen (1997); the adopted rotation velocity V_c is 220 km s^{-1} . The remaining inputs were the same adopted in García-Berro et al. (1998). In order to reproduce accurately the properties of the real catalog both MC simulations were required to meet the additional set of criteria: $\delta \geq 0^\circ$, $8.5 \leq M_V \leq 16.5$, $\mu \leq 4.1'' \text{ yr}^{-1}$ and $0.006'' \leq \pi \leq 0.376''$, which are derived from the subset of 232 white dwarfs which have all the determinations. An added value of the above described procedure of mixing tracer and real stars is that in this way we can check the accuracy of the classification algorithm and the quality of the MC simulations.

The simulated samples mimic fairly well the observational sample as can be seen in figure 1, where the results of the MC simulations for the disk and the halo are compared with the observational sample in the reduced proper motion-color diagram. As can be seen in this diagram the two simulated samples are easily visualized. Similar diagrams can be produced for each pair of variables and the results of the MC simulations compare equally well with the real data.

We have run the public domain neural network software SOM_PAK (available at http://www.cis.hut.fi/nsrc/som_pak/) with a catalog constructed as described above. The SOM of the input catalog, after three passes over the entire sample and with a grid of 5×5 nodes is shown in figure 2. The groups have been assigned either to halo (“H”) or disk (“D”) populations if the percentage of tracer stars of one of the populations was larger than 70%. In the groups labeled as “I” (intermediate population) neither the halo nor the disk tracers were in excess of this recognition percentage. As can be seen in figure 2 all the halo groups are close neighbors and, furthermore, the intermediate population

groups are surrounded by halo and disk groups. A good measure of the overall quality of the classification scheme can be obtained by checking how many of the synthetic stars are missclassified. This results in the following confusion matrix

$$C = \begin{pmatrix} 0.98 & 0.03 \\ 0.02 & 0.97 \end{pmatrix}$$

where the matrix element C_{11} indicates the percentage of disk tracers classified in disk groups, C_{21} is the percentage of disk tracers missclassified in halo groups, and so on. This matrix is very close to unity, and thus the classification seems to be secure. More confidence in this classification comes from the fact that the vast majority of old disk white dwarfs in the sample of Liebert, Dahn & Monet (1988) are in the groups (0,2) and (2,1) which are labeled as intermediate population. Moreover, LHS 56, LHS 147 and LHS 291 belong to the group (1,0) which clearly is a halo group, and LHS 2984 belongs to the group (0,0), all these objects were already identified as halo members by Liebert et al. (1989), and were used to build their halo white dwarf luminosity function. The only object of the sample of Liebert et al. (1989) missclassified is LHS 282 which is classified in the group (0,2), which is intermediate population. All this evidence points in the same direction: the classification is correct. The percentages of halo tracers in the groups labeled as halo can be found for each of the halo groups in figure 2. Since all of them are larger than 80% all these groups can be securely labeled as halo. However, and for the sake of reliability we have only identified as halo candidates those white dwarfs belonging to groups which do not have a disk neighbor, namely (0,0), (0,1), (1,0) and (2,0). One interesting property of these white dwarfs is that all of them have $M_V \leq 14$ and only 4 have proper motions in excess of $1.0'' \text{ yr}^{-1}$, being the average $\langle \mu \rangle = 0.87'' \text{ yr}^{-1}$. However, most of them have $\pi \leq 0.03''$, and are clustered around $\pi \sim 0.01''$, leading to tangential velocities in excess of 200 km s^{-1} for 11 of our candidates. Only one candidate has a tangential velocity smaller than 100 km s^{-1} . Therefore the

detected population is intrinsically bright and distant. The halo white dwarf candidates detected here can be found in table 1.

EDITOR: PLACE TABLE 1 HERE.

We have used the $1/V_{\text{max}}$ method to derive a luminosity function of halo white dwarfs with the candidates found so far. The adopted criteria for deriving such a luminosity function were those of Oswalt et al. (1996). The result is shown in figure 3 as solid triangles. The error bars have been computed as in Liebert et al. (1988). The number of objects in each luminosity bin is shown on top of the corresponding error bar. The value of $\langle V/V_{\text{max}} \rangle = 0.115$ indicates that the sample is not complete. Thus, this halo white dwarf luminosity function should be considered as preliminary although represents a considerable improvement over that of Liebert et al. (1989). According to the previous discussion, the most obvious feature of this luminosity function is the detection of halo candidates for the brightest luminosity bins. The faintest halo candidate found by Liebert et al. (1989), LHS 282, has been classified as intermediate population and, therefore, the faintest bin in their luminosity function is absent in ours. Some other small differences for the faintest bins are due to the binning procedure and to the bolometric magnitudes assigned to individual objects — we have used the bolometric corrections of Bergeron et al. (1995), whereas Liebert et al. (1989) used blackbody bolometric corrections.

For comparison purposes in figure 3 we also show two theoretical luminosity functions computed with a standard IMF. The adopted ages were in both cases 14 Gyr, and the durations of the bursts were 1 Gyr. The solid line corresponds to a luminosity function computed with a H envelope (Wood 1995) whereas the dotted line shows the luminosity function obtained using a He dominated envelope (Wood & Winget 1989). The method used to compute both luminosity functions is that of Isern et al. (1998), where the rest

of the details of the adopted inputs can be found. Also shown in figure 3 is the detection limit of very faint white dwarfs (open triangles) of Liebert et al. (1988). Both luminosity functions have been normalized to the bin with the smallest error bars and present a reasonable agreement with both sets of observational data.

3. Conclusions

We have shown that an artificial intelligence algorithm is able to classify the catalog of spectroscopically identified white dwarfs and ultimately detect several potential halo white dwarfs. Some of these white dwarfs were already proposed as halo objects by Liebert et al. (1989). We have found as well that our halo candidates are bright and distant, and that most of them have large tangential velocities. Using the $1/V_{\text{max}}$ method we have computed a preliminary luminosity function of halo white dwarfs. We have found a value of $\langle V/V_{\text{max}} \rangle = 0.115$ which indicates that our luminosity function is still uncomplete. However, this luminosity function largely improves the previous one by Liebert et al. (1989). We have also compared this luminosity function with the theoretical predictions, and we have found a fair agreement with the luminosity functions computed with a standard IMF.

Acknowledgements This work has been supported by DGICYT grants PB94-0111, PB94-0827-C02-02, and by the CIRIT grant GRC94-8001.

REFERENCES

- Adams, F., Laughlin, G., 1996, ApJ, 468, 586.
- Alcock, C., Allsman, R.A., Alves, D., Axelrod, T.S., Becker, A.C., Bennett, D.P., Cook, K.H., Freeman, K.C., Griest, K., Guern, J., Lehner, M.J., Mashall, S.L., Peterson, B.A., Pratt, M.R., Quinn, P.J., Rodgers, A.W., Stubbs, C.W., Sutherland, W., Welch, D.L., 1997, ApJ, 486, 697
- Bazell, D, Peng, Y., 1998, ApJS, 116, 47
- Bergeron, P., Wesemael, F., Beauchamp, A., 1995, PASP, 107, 1047
- Chabrier, G., Ségretrain, L., Méra, D., 1996, ApJ, 468, L21.
- Flynn, C., Gould, A., Bahcall, J.N., 1996, ApJ, 466, L55
- García-Berro, E., Torres, S., 1997, in “*White Dwarfs*”, Eds. J. Isern, M. Hernanz & E. García-Berro (Dordrecht: Kluwer), 97
- García-Berro, E., Torres, S., Isern, J., Burkert, A., 1998, MNRAS, submitted
- Hernández-Pajares, M., Floris, J., 1994, MNRAS, 268, 444
- Isern, J., García-Berro, E., Hernanz, M., Mochkovitch, R., Torres, S., 1998, ApJ, 503, 239
- Kohonen, T., 1990, Proc. IEEE, 78, No. 9, 1464
- Kohonen, T., 1997, “*Self-Organizing Maps*”, Springer Series in Information Sciences, Vol. 30 (Berlin: Springer)
- Liebert, J., Dahn, C.C., Monet, D.G., 1988, ApJ, 332, 891
- Liebert, J., Dahn, C.C., Monet, D.G., 1989, in “*White Dwarfs*”, Ed. G. Wegner (Berlin: Springer), 15
- Marković, D., Sommer-Larsen, J., 1997, MNRAS, 288, 733
- McCook, G.P., Sion, E.M., 1987, ApJS, 65, 603

- Méndez, R.A., Minnitti, D., De Marchi, G., Baker, A., Couch, W.J., 1996, MNRAS, 283, 666.
- Mochkovitch, R., García-Berro, E., Hernanz, M., Isern, J., Panis, J.F., 1990, A&A, 233, 456
- Naim, A., Lahav, O., Sodre, L., Storrie-Lombardi, M.C., 1995, MNRAS, 275, 567
- Oswalt, T.D., Smith, J.A., Wood, M.A., Hintzen, P., 1996, Nature, 382, 692
- Salpeter E.E., 1961, ApJ, 134, 669
- Serra-Ricart, M., Aparicio, A., Garrido, L., Gaitan, V., 1996, ApJ, 462, 221
- Tamanaha, C.M., Silk, J., Wood, M.A., Winget, D.E., 1990, ApJ, 358, 164
- Wood, M.A., 1995, in “*White Dwarfs*”, Eds. D. Koester & K. Werner (Berlin: Springer), 41
- Wood, M.A., Winget, D.E., 1989, in “*White Dwarfs*”, Ed. G. Wegner (Berlin: Springer), 282

Table 1. Halo white dwarf candidates identified using the neural network algorithm, along with their corresponding group and properties. The stars already identified in Liebert et al. (1989) are marked with an asterisk.

Name	Group	M_V	μ (″ yr ⁻¹)	π (″)	$B - V$	Sp. Type
LHS 2984*	(0,0)	11.62	0.930	0.015	0.03	DA
LHS 3007	(0,0)	13.06	0.636	0.028	0.29	DA
G 028-027	(0,0)	12.41	0.281	0.003	0.03	DQ
G 098-018	(0,0)	11.81	0.426	0.003	0.38	DA
G 138-056	(0,0)	13.34	0.692	0.006	0.37	DA
G 184-012	(0,0)	13.18	0.427	0.017	0.26	DC
LP 640-069	(0,0)	12.75	0.284	0.009	0.29	DA
LHS 56*	(0,1)	13.51	3.599	0.069	0.36	DA
LHS 147*	(0,1)	13.64	2.474	0.016	0.40	DC
LHS 151	(0,1)	13.46	1.142	0.053	0.33	DA
LHS 291*	(0,1)	13.39	1.765	0.012	0.11	DQ
LHS 529	(0,1)	13.94	1.281	0.046	0.64	DA
LHS 1927	(1,0)	11.41	0.661	0.009	0.11	DA
G 038-004	(1,0)	12.31	0.428	0.010	0.17	DA
LHS 3146	(2,0)	11.88	0.579	0.024	0.17	DA
G 021-015	(2,0)	11.54	0.390	0.015	0.05	DA
G 035-026	(2,0)	11.12	0.335	0.007	-0.14	DA
G 128-072	(2,0)	12.53	0.457	0.025	0.21	DA
G 271-106	(2,0)	11.77	0.396	0.014	0.18	DA
GR 363	(2,0)	11.39	0.133	0.003	-0.03	DA




Fig. 1.— Reduced proper motion-color diagram for the MC simulations of the disk (solid dots) and the halo (solid squares) and of the observational sub-sample (open triangles).




Fig. 2.— Self-organizing map of the sample of white dwarfs, see text for details. The group (0,0) is located in the lower left corner of the diagram and the group (4,4) is located in the upper right corner. As a rule of thumb H increases from right to left and M_V decreases downwards in the diagram.




Fig. 3.— White dwarf luminosity function obtained using the $1/V_{\max}$ method with the halo white dwarf candidates found in this work — solid triangles — and the detection limit of Liebert et al. (1988) — open triangles. The solid and the dotted lines are theoretical white dwarf luminosity functions obtained assuming a standard IMF and thick H and He dominated envelopes, respectively.